



Ragtime

MAPP 13259

RAGTIME is for spot treatment of ragwort and other perennial weeds in grass paddocks and lawns.

A ready to use soluble concentrate containing 4.82g/L 2,4-D and 1.68 g/L dicamba as the dimethylamine salt.

Batch: XXXX

FOR USE ONLY AS A HOME AND GARDEN WEEDKILLER

TO AVOID RISK TO MAN AND THE ENVIRONMENT, COMPLY WITH THE INSTRUCTIONS FOR USE.

Contains 2,4-D. May produce an allergic reaction.

Contents:

5L

Treats:

5L treats 200 square meters

PROTECT FROM FROST

Nufarm UK Limited
Crabtree Manorway North
Belvedere
Kent
DA17 6BQ
Fax: +44 (0) 208 319 7200

Technical Helpline telephone number: 0208 319 7232

24-hour emergency telephone number: 0208 319 7289

INSTRUCTIONS FOR USE

When to use

RAGTIME is a herbicide for the control of common ragwort and other perennial weeds in grass paddocks and lawns.

Ragwort


Spray when the majority of plants are in the rosette stage and growing vigorously in the autumn or spring but before the flower spines start to grow.

Fields to be grazed by animals should be treated in the spring. Ragwort sprayed with a herbicide will lose its natural repellent smell and becomes more attractive and palatable to grazing animals. It is vital no grazing animals are returned to the sprayed area until the ragwort has died and the dead leaves are fully decayed. This can take from 4-8 weeks after application.

Common ragwort will require spraying over a minimum of 2-years for complete control.

If rain falls shortly after application, the effect of RAGTIME may be reduced.

The correct growth stage for ragwort is illustrated below:



7cm

DO NOT mow or roll grass for four days before or after application. Clippings from the first four mowings after treatment must be composted for at least 6 months before use.

DO NOT treat grass suffering from drought, disease or other adverse factors.

DO NOT apply when grass is flowering.

DO NOT apply more than two treatments per year

WHERE TO USE

For use on grass paddocks and lawns

DO NOT treat newly established grass less than 1 year old.

HOW TO USE

RAGTIME is a 'ready to use' herbicide and requires no mixing. Turn the nozzle to 'spray' position. Hold the sprayer approximately 30cm from the ground, aiming at ragwort weeds and spray. Apply as a medium spray to just before run-off occurs.

[(For use with the 5L sprayer) Follow container assembly instructions. Hold the sprayer approximately 30cm from the ground, aiming at ragwort weeds and spray. Apply as a medium spray to just before run-off occurs.]

REFILLS

When using the refill, carefully pour into the sprayer ensuring no spillages occur. If it comes into contact with skin, wash off with plenty of water

Keep spray away from non-target plants

SAFETY INSTRUCTIONS

Protection during use

AVOID ALL CONTACT BY MOUTH

WASH HANDS AFTER USE

Environmental protection

Do not contaminate water with product or its container

TOXIC TO AQUATIC ORGANISMS, may cause long term adverse effects in the aquatic environment.

Do not contaminate surface waters or ditches

HARMFUL TO LIVESTOCK. Keep livestock out of treat areas for at least 14 days and until foliage of any poisonous weeds such as ragwort has died and become unpalatable.

Storage and disposal

KEEP PRODUCT AWAY FROM CHILDREN AND PETS

KEEP AWAY FROM FOOD AND DRINK

KEEP PRODUCT IN ORIGINAL CONTAINER, tightly closed, in a safe place

EMPTY CONTAINER COMPLETELY and dispose of safely

Dispose of unused product in its container at a household waster site (civic amenity site). Contact your local council (local authority) for details.

Dispose of empty container in the dustbin.