

---

# Holster

## Product Information

---

### GENERAL INFORMATION

Holster is a turf herbicide that has been specifically developed for the control of problem weeds in amenity turf (including fine turf).

Holster provides outstanding control of a range of broad-leaved weeds including Yellow Suckling Clover, Slender Speedwell, Daisy and Common Mouse Ear.

Holster contains three powerful active ingredients – fluroxypyr, 2,4-D and dicamba.

Fluroxypyr is a synthetic plant auxin and belongs to a chemical group called pyridine carboxylic acids (PCA's). Fluroxypyr is active at very low dose rates and is readily translocated through the target weeds.

Dicamba and 2,4-D also work by interrupting plant hormone activity in susceptible weed species.

Weeds treated with Holster show symptoms such as twisted leaf expansion and deformed growth - ultimately leading to the death of the plant.

---

### APPLICATION

Apply when the weeds are actively growing (normally between April and September) and when the soil is moist. Best results will be obtained by application in spring or early summer before weeds begin to flower.

New turf may be treated in the spring provided that at least two months have elapsed between sowing and treatment.

Avoid overlap or accidental overdosing – particularly on newly sown turf.

Avoid mowing for three days before and after spraying.

### RATES OF USE

<b>Method of Application</b>	<b>Rate of Holster</b>	<b>Water Rate</b>	<b>Area Treated</b>
Powered Sprayer	2.0 litres	200 – 1000 litres	1 hectare
Knapsack Sprayer	20 ml	10 litres	100 sq m
Walkover	100 ml	25 litres (467 L/ha)	525 sq m
Groundsman Walkover Professional	135 ml	25 litres (374 L/ha)	668 sq m

Holster should be applied as a medium quality spray.

## WEED CONTROL

The following weeds are controlled by Holster:

Cat's ear	Daisy	Slender Speedwell
Common Chickweed	Dandelion	White Clover
Common Mouse-ear	Mouse-ear Hawkweed	Yarrow
Common Sorrel	Plantain spp	Yellow Suckling Clover
Creeping Buttercup	Selfheal	



---

## IMPORTANT NOTES

Do not apply if rainfall is expected within 4 hours of application.

Avoid drift on to non target plants as very small quantities of Holster may cause severe damage to desirable plants.

Do not spray if turf is wet.

Do not spray in drought conditions or if turf is under stress from frost, waterlogging, trace element deficiency, pest or disease attack.

Repeat application 8 – 10 weeks after first application if necessary.

## PACK SIZE

2 Litres – treats 1.0 hectare

**Always read the label. Use pesticides safety.**

Holster is a registered trademark of Barclay Chemicals Manufacturing Ltd