

Strengthen your crop

syngenta®  
Biologicals

BIOSTIMULANT

# Maxicrop® Xtra-Fol

**Maxicrop Xtra-Fol is a liquid fertiliser containing NPK, Sulphur, trace elements, plant extracts and Maxicrop seaweed extract**



- It can be safely used on a wide range of agricultural and horticultural crops, to help maintain plant development during periods of rapid growth or stress (when a plant's ability to take up soil nutrients may be limited).
- The trace element package provides a balanced supplement for strong growth, though it should be noted that it will not correct severe micro nutrient deficiencies.
- The seaweed extract in Maxicrop Xtra-Fol stimulates root growth, contributes to chlorophyll retention and helps plants cope better with environmental and abiotic stress factors. The net effect is to produce strong crops for optimum yield and quality.

## MODE OF APPLICATION



SOIL/ROOT



FOLIAR



10L

Powered by



GEAPOW®

# Maxicrop® Xtra-Fol

## COMPOSITION

### COMPOUND FLUID FERTILISER (w/w)

Total Nitrogen (N) 8% - Phosphorus Pentoxide soluble in water (P<sub>2</sub>O<sub>5</sub>) 8% (P=3.5%)

Total Potassium Oxide (K<sub>2</sub>O) 7% (K=5.8%) - Sulphur (SO<sub>3</sub>) 2.5%

### TRACE ELEMENTS

Boron (B) 0.01%; water soluble Copper (Cu) 0.002%; Copper (Cu) chelated by EDTA 0.002%; water soluble Iron (Fe) 0.07%; Iron (Fe) chelated by EDTA 0.07%; water soluble Manganese (Mn) 0.017%; Manganese (Mn) chelated by EDTA 0.017%; water soluble Zinc (Zn) 0.01%; Zinc (Zn) chelated by EDTA 0.01%, Molybdenum (Mo) 0.0012%.

## DIRECTIONS

**FOLIAR SPRAYER:** Apply at 3-5 litres per ha in 200-600 litres of water/ha.

Apply when enough leaf cover is available, at 10-14 day intervals or as required.

**KNAPSACK SPRAYER:** Apply 100 ml in 10 litres of water at 10-14 day intervals or as required, thoroughly wetting the foliage.

**SOIL DRENCH/FERTIGATION:** Mix 3ml /litre of water (1:300 dilution). Apply to the root zone at 7-14 day intervals, or as required.

## WARNINGS

EUH208 Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

**HANDLING:** Wash hands after use.

**STORAGE:** Keep in original container and reseal after use. Store out of direct sunlight, above 0°C and below 30°C. Keep out of reach of children and pets.

**MIXING:** Shake before use. Once diluted, use straight away and flush equipment and pipework with water after use.

**TANK MIX COMPATIBILITIES:** Due to the wide range of different spray products available, it is impossible to test all combinations and we cannot accept any liability for any loss or damage. If in doubt, carry out a bucket test to check compatibility. Do not apply under adverse weather conditions, including frost, heavy rainfall, drying winds, extreme heat and extreme drought.

### Syngenta UK Limited

Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY

Tel +44 (0) 1223 883400

L1143426 GBRI/09D

### In case of toxic or transport emergency ring

**+44 (0)1484 538444 any time**

Maxicrop® is a registered trademark of SBM Développement.