



GUARD 2000

Contains 10% w/w styrene-butadiene co-polymer.

Authorisation No. ADJ 0652

FOR USE ONLY AS AGRICULTURAL/HORTICULTURAL/FORESTRY STICKER/EXTENDER

STATUTORY CONDITIONS RELATING TO USE:
For use in tank mixes with approved pesticides.
Maximum Concentration:

Maximum Number of Treatments:

Latest Timing:

On all edible crops:
With all approved pesticides applied at half
or less than half of approved rate.
With approved potato blight fungicides
With approved pesticides on other crops up to and including the growth stages given below.

0.1% of water volume.

Follow Statutory conditions of use of the pesticide.

Follow Statutory conditions of use of the pesticide.

Follow Statutory conditions of use of the pesticide.

Follow Statutory conditions of use of the pesticide.

SAFETY PRECAUTIONS

Causes serious eye damage.

Danger

Wear protective
gloves/protective
clothing/eye
protection/face protection.
IF IN EYES:

Rinse cautiously with water for
several minutes. Remove
contact lenses, if present and
easy to do. Continue rinsing.

Contains:

POLYOXYETHYLENE (7)TRIDECYL ETHER

The (COSHH) Control of Substances Hazardous to
Health Regulations may apply to the use of this product at
work.

OPERATOR PROTECTION

Follow all operator protection instructions for the partner
pesticide. Where the personal protective equipment specified
on this label is different to that specified on the pesticide
label the highest level of protective equipment must be worn.

OTHER SPECIFIC RESTRICTION

Do not use Guard 2000 with any pesticide which is to be
applied in or near water.

PROTECT FROM FROST

Made in UK

Batch No: See neck of the bottle



CROP	LATEST TIME OF APPLICATION
Broad, dwarf French, field and runner beans	First pod set.
Vining and combining peas	Flat pod (non-edible pods only)
Edible podded peas	Pod set
Oilseed rape	10% potential pods
Linseed	10% capsules formed
Sugar beet	6 fully expanded leaves
Cereals	GS52 (1/4 inflorescence)
Maize	Tip of tassel visible
Brussels sprouts	Lateral buds beginning to develop
Potatoes (except with approved potato blight fungicides – see above)	Tuber initiation
Cauliflower	Heads beginning to form (width up to 1cm)
Cucumber	First fruit set
Tomato	First fruit set on truss
Strawberries	First fruit development
Raspberries	First fruit set
Blackcurrants	First visible green fruitlet
Apples/pears	Fruits 5 - 10 mm
Cherry/plum	First fruit set
Apricot, almond, peach	First fruit set
Wine and table grapes	First fruit set
Hops	Cone set
Lettuce	Varieties forming heads – heads beginning to form Varieties not forming heads – when leaf rosette has reached 30% of typical diameter

These latest timings do not apply when using Guard 2000 with crop nutrition products and/or
seaweed-based plant growth stimulants.

Contents: 2.5 litres

15096937
10102016

Headland Agrochemicals Ltd., Rectors Lane, Pentre, Flintshire, CH5 2DH, UK.

Tel: + 44 (0) 1244 537370 Fax: + 44 (0) 1244 532097

E-mail: flintshire.enquiry@fmc.com www.headlandcroppnutrition.com

FMC

GUARD 2000

Authorisation No. ADJ 0652

Guard 2000 is a sticker/extender for use in agriculture and horticulture and is specially formulated for tank-mixing with a wide range of foliar-applied pesticides, with crop nutrition products and with seaweed based plant growth stimulants. Guard 2000 is an organic copolymer/ biodegradable surfactant blend and is designed to provide:

Rainfastness

Improved foliar cover

Improved foliar retention of spray

Protection against irrigation wash-off

Protection from u-v decomposition

Protection against evaporation loss

Guard 2000 may also be used to reduce drift where this is a concern.

USE SITUATIONS

Guard 2000 should be used in situations where chemical or micro-nutrient performance can be adversely affected by unfavourable weather, crop or growing conditions.

DIRECTIONS FOR USE

Guard 2000 may be used with approved pesticides in accordance with the Statutory Conditions of use and with foliar-applied micro-nutrients and seaweed-based plant growth stimulants. The Statutory Conditions Relating to Use of any partner pesticide must also be complied with.

Use Guard 2000 at 0.1% of total spray volume (100 ml per 100 litres of water or pro rata).

MIXING

Prepare the pesticide spray mixture in accordance with the manufacturers instructions. Add the required amount of Guard 2000 to the spray mix as the last operation. Maintain agitation

whilst mixing, travelling and spraying.

Clean spray equipment thoroughly after use.

COMPATIBILITY

Guard 2000 is compatible with most foliar-applied pesticides. Consult your distributor for information. Alternatively full compatibility information may be found at www.headlandcroppnutrition.com. Note: Do not mix Guard 2000 with formulations of bifenox.

Guard 2000's superior adhesive qualities may interfere with the vapour phase activity of pirimicarb formulations. Mixtures with these are not advised.

STORAGE

Protect from frost. Store above 5°C.

Keep in cool, well ventilated suitable pesticide store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

PRECAUTIONS

FOLLOW ALL OPERATOR PROTECTION INSTRUCTIONS FOR THE PESTICIDE.

Where the personal protective equipment specified on the adjuvant label is different to that specified on the pesticide label, the highest level of protective equipment must be worn. DO NOT USE with any pesticide which is to be applied in or near water.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

15096936
10102016

Headland Agrochemicals Ltd., Rectors Lane, Pentre, Flintshire, CH5 2DH, UK.
Tel: + 44 (0) 1244 537370 Fax: + 44 (0) 1244 532097
E-mail: flintshire.enquiry@fmc.com www.headlandcroppnutrition.com

FMC