

Kurtail Evo



MAPP 19020

A soluble concentrate formulation containing 160 g/l 2,4-D and 240 g/l glyphosate.

- Controls annual and perennial weeds in stubbles, grassland (destruction), green cover on land not being used for crop production, orchards, amenity grassland (destruction), hard surfaces (railway ballast only) and amenity vegetation.

Controls weeds which are difficult to kill with straight glyphosate.

LERAP category: B

Storage: Protect from frost

Kurtail Evo

Contains 160 g/l 2,4-D and 240 g/l glyphosate.



WARNING

May cause an allergic skin reaction. Causes serious eye irritation. Very toxic to aquatic life with long lasting effects.

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.

Avoid breathing vapours or spray. Do not eat, drink or smoke when using this product.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

DANGER.

To avoid risks to human health and the environment, comply with the instructions for use.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

PROTECT FROM FROST

Progreen Weed Control Solutions Ltd, Unit 7 Spalding Road Business Park, Bourne, PE10 9LF
Telephone Number: 0800 032 6262

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Maximum individual dose	Maximum no. of applications	Latest time of application
All edible crops (stubble), all non-edible crops (stubble)	5.0 L/ha	One	Five days before drilling or planting of the following crop
Grassland			Five days before harvest, grazing or drilling
Green cover on land not being used for crop production			Five days before drilling or planting of the following crop
Apple (around), pear (around)			After harvest but before green cluster stage
Amenity grassland			Five days before drilling or planting of the following crop
Hard surfaces (railway ballast only)			-
Amenity vegetation			

Other specific restrictions:

- Do not use under polythene or glass
- Application of lime, fertiliser, farmyard manure and pesticides should be delayed until five days after application of KYLEO.
- After application large quantities of decaying foliage, stolons, roots or rhizomes should be dispersed or buried by thorough cultivation before crop drilling.
- Do not sue in or alongside hedges.
- Do not apply by hand-held equipment.
- Windfall (fruit fall to ground) must not be used as food or feeding stuff.
- This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with the 'Local Environment Risk Assessment for Pesticide Horizontal Boom Sprayers' booklet available from the HSE Chemicals Regulation Directorate's website or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS**Operator Protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental Protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with the product or its container.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within five metres of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone or within one metre of the top of a ditch which is dry at the time of application. Direct spray away from water.



This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with the 'Local Environment Risk Assessment for Pesticides Horizontal Boom Sprayers' booklet available from the HSE Chemicals Regulation Directorate's website or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Do not contaminate water with the product or its container.

(Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads.)

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area. To protect aquatic organisms/non-target plants do not apply on impermeable surfaces such as asphalt, concrete, cobblestones and other situations with a high risk of run-off.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, and dispose of safely.

This material and its container must be disposed of in a safe way.

DO NOT RE-USE CONTAINER for any purpose.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

WARNINGS

EXTREME CARE SHOULD BE TAKEN TO AVOID SPRAY DRIFT AS THIS CAN SEVERLY DAMAGE OR DESTROY NEIGHBOURING CROPS AND PLANTS.

RESTRICTIONS

DO NOT spray in windy conditions as drift onto desired crops or vegetation could severely damage or destroy them.

DO NOT spray onto weeds which are naturally senescent, or where growth is impaired by drought, high temperatures, a covering of dust, flooding or frost at, or immediately after application, otherwise poor control may result.

Do not use under polythene or glass.

After application large quantities of decaying foliage, stolons, roots or rhizomes should be dispersed or buried by thorough cultivation before crop.

Application of lime, fertiliser, farmyard manure and pesticides should be delayed until five days after application of KYLEO.

Do not use in or alongside hedges.

Windfall (fruit fall to ground) must not be used as food or feeding stuff.

Do not apply by hand-held equipment.

WEEDS CONTROLLED

Susceptible

Bindweed, black	Knotgrass
Bindweed, field	Mayweed, spp
Cleavers	Meadow grass, annual
Couch, common	Medick, black
Dandelion	Ryegrass
Deadnettle, red	Shepherd's purse
Dock, broadleaved	Speedwell spp
Fat hen	Thistle, creeping
Forget-me-not, field	Volunteer cereals
Groundsel	Volunteer oilseed rape

Annual weeds susceptible at 3.0 L/ha

Perennial weeds susceptible at 5.0 L/ha

Annual and perennial grass weeds susceptible at 5.0 L/ha

Broad mix of weeds with volunteer cereals susceptible at 5.0 L/ha

Annual grasses and broadleaved weeds should have at least 5cm of leaf, or two expanded true leaves respectively.

Perennial grasses must have a full emergence of healthy, green leaf. Common couch for example becomes susceptible at the onset of tillering and new rhizome growth commences which usually occurs when plants have four to five leaves, each with 10-15 cm of new growth.

KYLEO will not give an acceptable level of control of horsetails.

CROP SPECIFIC INFORMATION

ALL edible crops (stubble), all non-edible crops (stubble)

Maximum individual dose 5.0 L/ha

Maximum total dose 5.0 L/ha

Timing

Apply following harvest and before sowing/planting of the following crop. Do not cultivate immediately before or after spraying. Clear straw or cut it short and spread evenly. Treat couch grass when it has formed three to four leaves and it is actively growing.

Water volume 100-500 litres of water per hectare.

Grassland

Maximum individual dose 5.0 L/ha Maximum total dose 5.0 L/ha

Timing

Treat either before grazing/mowing when growth is 30-60 cm, not dense and lacking mature seeds, or re-growth after grazing/mowing. Treat couch grass when it has formed three to four leaves and is actively growing. For grassland renewal treatment and re-sowing in July and August have proved effective.

Water Volume 100 – 500 litres of water per hectare.

Green cover on land not being used for crop production e.g. Set Aside

Maximum individual dose 5.0 L/ha

Maximum total dose 5.0 L/ha

Timing

For perennial weed control allow at least 21 days before spraying in spring and allow five days before cultivating or drilling. For annual weed control allow 24 hours before cultivating. Treat couch grass when it has formed three to four leaves and is actively growing. Ensure that all management rules are followed prior to use on land taken out of production as part of a grant aided scheme.

Water volume 100 – 500 litres of water per hectare.

Apple (around), Pear (around)

Maximum individual dose 5.0 L/ha

Maximum total dose 5.0 L/ha

Timing In spring or summer OR after harvest

Application guidance

Trees must have established for three years before spraying.

Avoid contact with tree trunks and branches. Do not use any fallen fruit as food or foodstuffs. Do not spray during flowering. Remove root suckers and side shoots before spraying. Do not spray in temperatures above 25°C and in windy weather. On hot and dry days, spray in the evening.

Pre-harvest interval 30 days if sprayed in spring or summer.

Water volume 100 – 500 litres of water per hectare

Amenity grassland**Maximum individual dose** 5.0 L/ha**Maximum total dose** 5.0 L/ha**Timing**

Apply when weeds have fully grown. Treat couch grass when it has formed three to four leaves.

Water volume 100 – 500 litres of water per hectare**Hard surfaces**

Kurtail Evo may be used on railway track, railway sidings and other ballast areas of rail infrastructure, station areas and surrounds.

Maximum individual dose 5.0 L/ha**Maximum total dose** 5.0 L/ha**Timing**

Apply Kurtail Evo post-emergence of weeds at any time from March until the end of September, provided that the weeds are actively growing and have not begun to tiller. Apply when weeds have fully grown. Treat couch grass when it has formed three to four leaves. Avoid drift on to crops alongside railway lines.

Water volume 100 – 500 litres of water per hectare**Amenity vegetation**

Apply to areas of bare soil around ornamental plants or areas intended for ornamental planting. Take care to avoid any contact with ornamental plants as this can severely damage them.

Maximum individual dose 5.0 L/ha**Maximum total dose** 5.0 L/ha**Timing**

Apply when weeds have fully grown. Treat couch grass when it has formed three to four leaves.

Water volume 100 – 500 litres per hectare**Following crops**

Soil cultivation can be carried out as soon as first weeds show clear signs of Kurtail Evo taking effect.

For grassland renewal, ensure a fine firm seedbed for sowing into moist soil and that seed is sown at 1 – 2 cm depth. Following sowing the ground should be rolled.

Kurtail Evo may cause damage to succeeding crops. Observe the following periods of time between application and the cultivation of succeeding crops:

Crop	Days until sowing
------	-------------------

All edible crops (stubble), All non-edible crops (stubble)*	5
Maize Sorghum	14
Field bean Buckwheat Peas Fibre flax, linseed Alfalfa Phacelia Oilseed rape Mustard	28
Potatoes Sunflowers Sugar beet Vegetables	60

*Only cereals should be planted as a succeeding crop following application to all edible crops (stubble), all non-edible crops (stubble) with a five day sowing interval.

MIXING AND SPRAYING

Before use ensure that the spraying equipment has been thoroughly cleaned. Half-fill the spray tank with clean water. With the contents of the spray tank under re-circulation, add the measured quantities of Kurtail Evo through the filter. Top up the tank with water to the required level and maintain re-circulation until the tank is sprayed out.

Apply the recommended quantity of Kurtail Evo through a conventional hydraulic sprayer using a MEDIUM spray to cover the weed leaves evenly and thoroughly.

DO NOT leave the sprayer standing with chemical in it.

On completion of spraying, drain sprayer and wash out thoroughly using water with the addition of a suitable detergent.

WEATHER AND GROWING CONDITIONS

Apply to a dry crop when rain is not forecast for at least three hours. Optimum results are obtained when the weeds are actively growing under good soil and weather conditions. Reduced weed control may be obtained during drought or cold weather. If rain falls shortly after application, the effect of Kurtail Evo may be reduced.

COMPANY ADVISORY INFORMATION

This section is not part of the Product Label under REGULATION (EC) No 1107/2009. It provides additional advice on product use at the discretion of the applicant.

Certain weeds may develop resistance to Progreen Weed Control Solutions Ltd's products. Since such circumstances are beyond our control, Progreen Weed Control Solutions Ltd will be under no liability for any resulting loss or damage whatsoever.