Environmental Information Sheet LEYSTAR® MAPP 17921



An emulsifiable concentrate formulation containing 144 g/litre fluroxypyr-meptyl and 80 g/litre clopyralid, pyridine carboxylic acid herbicides, and 2.5 g/litre florasulam, a triazolopyrimidine herbicide, for use in grassland and forage maize.

Maximum individual dose: 2.0 l/ha (288 gas/ha fluroxypyr-meptyl, 160 gas/ha

clopyralid, 5.0 gas/ha florasulam)

Maximum number of applications: 1 per year

	Section	Profile
1.	WILDLIFE	LEYSTAR is not classified as "Harmful to game, wild birds and animals".
	Mammals and Birds	No risk management is necessary. LEYSTAR has low toxicity to mammals and low to moderate toxicity to birds. The risk to birds and mammals living and feeding in treated areas is low.
		"Livestock must be kept out of treated areas for at least 7 days following treatment. Animals should be excluded from treated areas until any ragwort has completely recovered or died and there is no visible sign of the dead weed."
2.	BEES	No risk management necessary. LEYSTAR has a low risk to honeybees and there is no requirement to avoid application of the product when bees may be foraging on flowering weeds.
3.	NON TARGET INSECTS AND OTHER ARTHROPODS	No risk management necessary. LEYSTAR poses a low risk to a range of arthropod species commonly found in and around treated fields, such as predatory mite (e.g. <i>Typhlodromus pyri</i>), aphid parasitoid wasp (<i>Aphidius rhopalosiphi</i>) and lacewing (<i>Chrysoperla carnea</i>).
4.	AQUATIC LIFE	LEYSTAR is classified "Very toxic to aquatic life with long lasting effects".
		LEYSTAR has a high toxicity to fish, <i>Daphnia</i> (water fleas) and algae and is extremely toxic to the aquatic plant <i>Lemna</i> . Care must be taken to avoid contamination of water with the product or its container. LEYSTAR can be used safely providing care is taken to minimise spray drift reaching surface waters.
		"DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water."
		Buffer zones may be reduced (see LERAP Guidelines).

[®]Trademark of the Dow Chemical Company (Dow) or an affiliate company of Dow

EIS Version 1 12 July 2017

Environmental Information Sheet LEYSTAR® MAPP 17921



Section		Profile
5.	SOIL and GROUNDWATER	The active substances are impersistent or of low persistence and breakdown products are impersistent to very persistent. All are of low to high mobility. However, when LEYSTAR is used as recommended there is a low risk of groundwater contamination.
	Earthworms	LEYSTAR has a moderate toxicity to earthworms. The risk to earthworms is low and no risk management is necessary.
	Soil Micro-organisms	No risk management necessary. LEYSTAR at the recommended rates has no significant effects on soil microbial respiration and nitrogen transformations. The risk to soil microbial activity is therefore low.
6.	NON-TARGET PLANTS	LEYSTAR is a post emergence herbicide active on a range of broad leaf weeds, consequently care should be taken to avoid spray drift reaching hedgerows, conservation headlands and adjacent crops to avoid possible effects on wild plants and sensitive crops. "Extreme care must be taken to avoid spray drift onto non-crop plants outside the target area".

ALWAYS READ THE LABEL: USE PESTICIDES SAFELY

Care must be taken to minimise the risk of surface water contamination from farmyard and field sources.

For further information about the environmental profile of this product contact:-

Company: Dow AgroSciences Limited

Address: CPC2 Capital Park, Fulbourn, Cambridge CB21 5XE

Telephone: 01462 457272
Fax: 01462 426605
e-mail: <u>ukhotline@dow.com</u>

24-hour Emergency Telephone Number; +44 (0)1553 761 251

This Environmental Information Sheet was prepared in accordance with CPA Guidance Notes Version 4.

© Copyright of Dow AgroSciences Ltd.

EIS Version 1 12 July 2017